
SOME INDICATORS OF A POSSIBLE METH LAB

Precursors (products used to make meth):

- More than a usual amount of precursors
- Lots of discarded precursor packages and containers
- Lots of precursors in the kitchen
- Car batteries inside the residence
- Propane tanks inside the residence
- Propane tanks with fittings that have turned blue (very dangerous!)
- Bottles or jars containing separated clear and opaque liquids
- Bottles or jars with rubber tubing attached
- Coffee filters containing a white pasty substance or a dark red sludge
- Glass cookware or frying pans containing a powdery residue

Property:

- Unusual and strong chemical odors, such as acetone or ammonia
- Windows blocked out by paint or dark bed sheets
- Electronic security devices, such as motion sensors or surveillance cameras
- Unexplainable explosions or fires

Visitors:

- Lots of nighttime visitors
- Lots of visitors who stay 5 or 10 minutes

Other:

- Renters who pay cash
- Persons exhibiting signs of meth use (see below)

SOME INDICATORS OF POSSIBLE METH USE

Increased heart rate

Dilated pupils

Bloodshot eyes

Dry mouth

Elevated body temperature

Excessive sweating

Chemical odor on the breath

Open sores or blemishes

False sense of confidence

Flushed or tense appearance

Excessive talking

Rapid or impaired speech

Aggressive behavior

Inability to sleep

Distorted sense of time

Loss of appetite

Loss of weight

Loss of memory

Loss of pleasure

Depression

Panic attacks

Paranoia

Hallucinations